

## CARROT SEED

Crop	Soil Depth	Depth to sodic layer	pH of top 15cm (H <sub>2</sub> O)	EC (top 15 cm)	Texture (top 15cm - % clay))	Drainage	Stoniness (top 15cm) Stone class	slope	Frost
W	>20		>=6	<1	>10%	Well drained, Mod well, Excess well	<10% <=2	<10%	At least 1 day where T <sub>min</sub> <-2°C at flowering (15 Dec – 15 Feb): Occurs in no years
S	>20		5.8-<6	1-1.7	>10%	Imperfectly drained	10-20% 3	<10%	At least 1 day where T <sub>min</sub> <-2°C at flowering (15 Dec – 15 Feb): Occurs <1 in 20 yrs
MS	15-20		5.5-<5.8	1.7-2.8	5-10%		20-35% 4	10-20%	At least 1 day where T <sub>min</sub> <-2°C at flowering (15 Dec – 15 Feb): Occurs 1/10 to 1/20yrs
U	<15		<5.5	>2.8	<5%	Poorly drained	>35% >4	>20%	At least 1 day where T <sub>min</sub> <-2°C at flowering (15 Dec – 15 Feb): Occurs >1 in 10

**Well suited (W):** Land having no significant limitations to sustained application of a given use, or only minor limitations that will not significantly reduce productivity or benefits and will not raise inputs above an acceptable level. Any risk of crop loss is inherently low or can be easily overcome with management practices that are easy and cheap to implement.

**Suitable (S):** Land having limitations which are moderately severe for sustained application of a given use; the limitations will reduce productivity or benefits and increase required inputs to the extent that the overall advantage to be gained from the use, although still attractive, will be appreciably inferior to that expected on Class S1 land. Risk of crop loss is moderately high or requires management practices that are difficult or costly to implement.

**Marginally Suitable (MS):** Land having limitations which are severe for sustained application of a given use and will so reduce productivity or benefits, or increase required inputs, that this expenditure will be only marginally justified. Risk of crop loss may be high.

**Unsuitable (U):** Land which has qualities that appear to preclude sustained use of the kind under consideration